

Q1 Gone fishing

5 Points

Q1.1

1 Point

You go fishing a very large number of times.

The number of fish you catch on a giving fishing trip is a random variable of some distribution.

Which distribution best approximates the total number of fish you catch (throughout all the fishing trips)?

- ☐ Binomial
- ☐ Geometric
- ☐ Poisson
- ☐ Exponential
- ☒ Normal

Explanation

The sum of a set of independent and identically distributed random variables approaches a normal distribution by CLT.

Q1.2**1 Point**

You catch an expected number of 1.5 fish per hour. You can catch a fish at any instant of time.

Which distribution best characterizes the number of fish you catch in one hour of fishing?

- ☐ Binomial
- ☐ Geometric
- ☒ Poisson
- ☐ Exponential
- ☐ Normal

Explanation

You're measuring the discrete number of fish you catch based on the rate at which fish-catching happens in a continuous time interval.

Q1.3**1 Point**

You catch an expected number of 1.5 fish per hour. You can catch a fish at any instant of time.

Which distribution best characterizes how the amount of time it takes to catch a fish after casting the line?

- ☐ Binomial
- ☐ Geometric
- ☐ Poisson
- ☒ Exponential
- ☐ Normal

Explanation

You're measuring the continuous duration of time until a fish is caught.

Q1.4

1 Point

Each day you fish for 2 hours. Assume that the numbers of fish you catch each day are independent.

Which distribution best characterizes the number of days until you catch more than 10 fish in one day?

- ☐ Binomial
- ☒ Geometric
- ☐ Poisson
- ☐ Exponential
- ☐ Normal

Explanation

The probabilities of catching more than 10 fish on each day are independently and identically distributed, and you're measuring the number of trials until success.

Q1.5

1 Point

Each day you fish for 2 hours. Assume that the numbers of fish you catch each day are independent.

Which distribution best characterizes the number of days you catch no fish over a week of fishing?

- ☒ Binomial
- ☐ Geometric
- ☐ Poisson
- ☐ Exponential
- ☐ Normal

Explanation

The probabilities of catching no fish on each day are independently and identically distributed, and you're measuring the total number of successful trials.

Q2 Normal distribution

2 Points

Let X be a Normal random variable with mean μ and variance σ^2 . Which of the following are true statements?

Q2.1

1 Point

$$\frac{X-\mu}{\sigma} \sim N(0, 1)$$

☒ True

☐ False

Explanation

$$\text{Let } Y = \frac{X-\mu}{\sigma}. \mathbb{P}[a \leq Y \leq b] = \mathbb{P}[a\sigma + \mu \leq X \leq b\sigma + \mu] = \frac{1}{\sqrt{2\pi\sigma^2}} \int_{a\sigma+\mu}^{b\sigma+\mu} e^{-(x-\mu)^2/(2\sigma^2)} dx = \frac{1}{\sqrt{2\pi}} \int_a^b e^{-y^2/2} dy.$$

Q2.2

1 Point

The sum of any two normal random variables is also normally distributed

☐ True

☒ False

Explanation

This is true if and only if the variables are independent. Proof under Corollary 21.1 in Note 21

Q2.3**0 Points**

The sum of n i.i.d. random variables follows a normal distribution as n approaches infinity.

☒ True

☐ False

Explanation

Let S_n be the sum of n i.i.d random variables each with mean μ , variance σ^2 . From CLT, $S_n \sim N(n\mu, n\sigma^2)$, which is normal.

Q3 CLT**1 Point**

Suppose $S_n = X_1 + \cdots + X_n$, where X_i 's are i.i.d random variables with $\mathbb{E}[X_i] = \mu$, $\text{Var}(X_i) = \sigma^2$. As $n \rightarrow \infty$, which of the following random variables converges to the standard normal distribution?

☐ $\frac{S_n}{n}$

☐ $\frac{S_n}{n} - \mu$

☒ $\frac{S_n - n\mu}{\sigma\sqrt{n}}$

☐ $\frac{S_n - n\mu}{\sigma n}$

Explanation

Note that by linearity of expectation, $\mathbb{E}\left[\frac{S_n - n\mu}{\sigma\sqrt{n}}\right] = \frac{1}{\sigma\sqrt{n}}(\mathbb{E}[S_n] - n\mu) = \frac{1}{\sigma\sqrt{n}}(\mathbb{E}[X_1] + \cdots + \mathbb{E}[X_n] - n\mu) = 0$, and since X_i 's are independent, $\text{Var}\left(\frac{S_n - n\mu}{\sigma\sqrt{n}}\right) = \frac{1}{\sigma^2 n} \text{Var}(S_n - n\mu) = \frac{1}{\sigma^2 n} \text{Var}(S_n) = \frac{1}{\sigma^2 n} (\text{Var}(X_1) + \cdots + \text{Var}(X_n)) = \frac{1}{\sigma^2 n} \cdot \sigma^2 n = 1$. By CLT, $\frac{S_n - n\mu}{\sigma\sqrt{n}}$ converges to $N(0, 1)$ as $n \rightarrow \infty$.